

7TITLE

Controlled release formulations containing an active ingredient, preferably melatonin, and the method of preparation

SUMMARY

6 This invention relates to new controlled release formulations containing an active ingredient, preferably melatonin, and the method of their preparation. Following their administration, the active ingredient, preferably melatonin, is released in with biphasic pattern and, in particular, rapidly during the first few minutes after its administration and later, slowly and gradually. Formulations obtained by such a process can be usefully employed in cases where controlled release of active ingredients contained in the formulation is needed, for example in disorders in the sleep/wake cycle and sleep disturbances, if they contain melatonin.

Description of the invention with the title:

Controlled release formulations containing an active ingredient, preferably melatonin, and the method for preparation.

In the name of: AMBROS PHARMA S.R.L.

Based in : MILAN

Designated inventor: Stankov Bojidar M.

* * * * *

FIELD OF APPLICATION

24 This invention relates to new controlled release formulations containing melatonin, characterized by the fact that the active ingredient is released quickly at first and slowly and gradually later.

Moreover, this invention relates to the use of those controlled release formulations as medicines, nutritional or health food supplements, for the treatment of sleep disturbances and to the new process of preparing controlled release formulations, especially those containing melatonin.

STATE OF THE ART

Melatonin (N-acetyl-5-methoxytryptamine) is an indolic compound widely present in Nature. It is produced by almost all living organisms, from algae to man. While in lower organisms melatonin could be part of the natural defence against oxidative stress, it is widely accepted that in mammals and man melatonin behaves as a *Zeitgeber* ("time giver"), that acting together with light (primary *Zeitgeber*), it is capable of synchronizing the endogenous biological clock with the phase of the prevalent photoperiod.

In this way melatonin controls the diurnal (circadian) rhythm of the organism, showing a strong synchronizing effect on the sleep/wake cycle. Because of its marked activity on the circadian clock, melatonin has been used in the treatment of syndromes related to the desynchronization of sleep/wake cycle, as, for example in cases of jet-lag , delayed sleep phase syndrome (DSPS), in blind persons, in the elderly and in insomnia in shift-workers and psycho-physiological insomnia.

Apart of its role in the control of the circadian rhythms, melatonin has also bland hypnotic effects which influence the induction, the depth and the quality of sleep. Indeed, melatonin is capable of improving the sleep micro-structure (CAP and CAP rate) CAP (Cycling Alternating Pattern) is the only objective parameter in establishing sleep quality.

Melatonin is also a strong regulator of body temperature and, since the lowering of body temperature is extremely important for the induction and quality of sleep, this gives further contribution of its effects on sleep.

In normal conditions, the night-time release of melatonin occurs with levels higher than 50-100 pg/ml that, during physiological sleep, last for about 6-7 hours. Under normal conditions melatonin very quickly reaches maximal physiological levels: within 30-45 minutes from the initiation of the nocturnal melatonin peak. This phenomenon has established, in research done on its effects on sleep, the choice of various doses (0.1 – 1 – 10 - 100 mg), a choice determined, on the other hand, by the short half-life of melatonin (20-40 minutes in various species, including man).

Because of the short half-life, the use of low doses melatonin (0.1 – 1 mg), that correspond to "physiological" levels of melatonin in the peripheral blood, proved to be

insufficient for the regulation of the sleep/wake cycle and sleep induction, because following a facilitation of the sleep induction phase, the subject shows the typical tendency to wake up, with high WASO (Wake After Sleep Onset) numerical values both, as the number of awakenings (NA) and as time spent awake after the beginning of the first sleep period.

6 The administration of high "pharmacological" doses of melatonin is to be as well avoided because of bioavailability of melatonin in the peripheral blood extending in the morning hours, when the melatonin levels physiologically have to be already very low.

In attempts to avoid this, melatonin has been administered in various "retard" formulations (oral, transdermal, transmucosal). However, a great disadvantage of those formulations is the relatively slow absorption of melatonin, often accompanied by low peripheral blood levels, with subsequent loss of the effect on sleep induction. Additionally, the transdermal patches show limited application, because of poor compliance and the danger of leaving (forgetting to take off) the patch in the morning, and therefore having high melatonin levels during daytime (residual release from the patch), when the melatonin levels have to be very low.

On the other hand, the controlled release forms (or pulsatile forms) for oral administration developed recently are very expensive, being prepared as a various number of tablets placed in the same capsule. Moreover, they are not suitable for the preparation of nutritional supplements, because of the use of synthetic retardants and excipients which are not allowed in nutrition and may have untoward effects.

24 SUMMARY

We have now found, and that is the subject of this invention, new formulations for the controlled release of melatonin able to "mimic" the physiological melatonin pattern in the peripheral blood.

The properties of the new formulations allow to vary the release parameters with the possibility of obtaining a defined biphasic pattern, similar to the natural one observed in
30 the organism during the nocturnal period of synthesis and endogenous release of melatonin.

These new formulations are prepared as a single tablet releasing its melatonin content in a controlled manner with the aim of obtaining a biphasic pattern of release. Both, *in vitro* and *in vivo*, there is initially an almost instant release (about 5-10 minutes after administration), of a pre-established percentage (for example between 25% and 30% of the total dose) followed by slow release of the rest of the melatonin contained in the tablet, with the start of the release about 30-45 minutes after administration, with a duration of 5-7 hours.

New slow-release formulations containing melatonin are, therefore, the subject of this invention. They are characterized by the fact they are made up of an internal slow-release, hard "nucleus", with a predetermined content of the active ingredient, and by a quick-release external layer ("cortex"), containing further predetermined doses of melatonin.

The content of melatonin can be between 0,1 and 100 mg inclusive, both in the internal nucleus and in the external "cortex", preferably between 1 and 3 mg in the internal nucleus and between 0,5 and 1,5 mg in the external "cortex".

The formulations according to this invention may be used both as pharmaceuticals and/or nutritional supplements, or functional foods, for the regulation of the sleep/wake cycle, or to substitute the melatonin levels decreasing with age or because of a pathological process.

A further object of this invention is a new process for the preparation of controlled release formulations, in particular those containing melatonin, characterized by the preparation of a nucleus containing an active ingredient, preferably melatonin, with a predetermined rigidity and by overcast of an external layer ("cortex") containing a further predetermined dose of active ingredient, preferably melatonin.

The procedure for the preparation of the said formulations is as follows:

- a) preparation of the delayed release nucleus containing the active ingredient, preferably melatonin;
- b) forming of the external layer ("cortex") containing a further dose of the main ingredient, preferably melatonin, under control;
- c) fixing of the external layer

DETAILED DESCRIPTION OF THE INVENTION

The use of formulations containing melatonin for the regulation of the sleep/wake cycle, for sleep induction and to substitute decreased melatonin levels has been known for a long time. In spite of this, up till now the exogenous administration of melatonin still presents serious problems connected with this compound's typical mode of action and its short half-life.

With the aim of obtaining a "physiological" effect it would however be necessary to achieve levels of melatonin able to induce sleep quickly and then to maintain the hormone levels such as to prolong the sleep period. Indeed, this is the only way to physiologically restore the melatonin rhythm and the equilibrium of the disrupted sleep/wake cycle.

As previously mentioned, none of the solutions adopted up until now have resolved this problem satisfactorily.

Moreover, the existing controlled release ("pulsatile-release") formulations, are not suitable for use as nutritional supplements or as foods. This is the preferred use within the embodiment of the present invention, knowing that melatonin shows no significant side-effects and is not a drug, but a natural compound.

With the aim of resolving previously unanswered problems, the Applicant has found new controlled release formulations for the administration of melatonin and a new and advantageous method for the preparation of controlled release formulations.

According to this invention, controlled release formulations are made in the form of a single tablet, characterized by a nucleus which is wrapped in an external layer ("cortex") both containing the active ingredient in predetermined doses.

To verify the effectiveness of the formulations according to the invention, we performed various tests where the doses of melatonin and the excipients used were determined according to the laboratory data and we compared the natural rhythm of melatonin in man, with pharmacokinetics data of the said formulations.

The aim of the invention was to obtain controlled release formulations in which about 100% of the active ingredient contained in the "cortex" is released in a period of between 5-10 minutes from its administration, and 80-90% of the active ingredient contained in the nucleus is released within four hours of its administration.

This biphasic release kinetics corresponds to two peaks in the concentration of melatonin, in the peripheral blood of man which occurs between thirty and one hundred and twenty minutes after its administration, with a bioavailability of about 5-6 hours following administration.

Further object of this invention therefore, is that of making new controlled release formulations available which permit to obtain maximum levels of melatonin in man equivalent to about 1,000 - 2,000 pg/ml, which correspond to very low pharmacological concentrations, but sufficient to obtain significant effects on the sleep/wake cycle, on sleep induction and structure and to substitute the decreased melatonin levels in cases, when necessary.

The formulations which are the subject of this invention are characterized by an internal nucleus and by an external layer ("cortex") in one single tablet, both containing the active ingredient, preferably melatonin, which releases the active ingredient biphasically. The melatonin content may be between 0.1 and 100 mg inclusive both in the internal nucleus and in the external "cortex". Preferably, it may be between 1 and 3 mg inclusive in the internal nucleus and between 0.5 and 1.5 mg in the external "cortex".

The excipients used in the preparation of the said formulations can be all those normally used in the field and suitable for the purposes of obtaining the desired controlled release, and in particular, all those permitted for use in the pharmaceutical field and especially in food supplement and/or nutritional field.

The method of preparing controlled release formulations containing the active ingredient, preferably melatonin, according to the invention is as follows:

- a) preparation of the delayed release nucleus containing the active ingredient, preferably melatonin;
- b) forming of the external layer ("cortex") containing a further dose of the main ingredient, preferably melatonin, under control;
- c) fixing of the external cover.

In particular, the new method of preparing controlled release formulations is described as follows:

- 1) Preparation of the mixture containing the active ingredient, in particular melatonin, for the formulation of a delayed release nucleus with the necessary excipients, in

particular volume excipients, gliding and lubricating excipients, binding and retardant excipients.

- 2) Preparation of the nucleus with a predetermined hardness
- 3) Preparation of the active ingredient solution, preferably melatonin, for the formation of the "cortex" of the tablets.

6 4) Application of the solution under pressure containing the active ingredient to the tablets to make a "cortex", with control of the quantity of the active ingredient by means of chemical analyses done on samples taken during the application. and at the end of the application

- 5) For this process all the excipients suitable for the purpose of this invention can be used. In particular for phase 1), dicalcium phosphate, mannitol, lactose, magnesium stearate, polyvinylpyrrolidon, and hydroxypropylmethylcellulose, viscosity 3,000-5,600 can be used, while for phase 3) the preferred choices are hydroxypropylmethylcellulose, lactose, ethyl alcohol and purified water.

Example 1

Method for preparing tablets containing melatonin according to the invention

Melatonin with a purity defined as HPLC >99.5% from the company HELSINN, (Biasca, Switzerland) was used. The rest of the ingredients were supplied by producers certified for the production and distribution of pharmaceuticals and/or food supplements. Hydroxypropylmethylcellulose (Methocel) with viscosity 3,000-5,600 from Dow Chemical Company was employed.

Preparation of the tablets

24 The tablets were produced in five batches: three pilot batches and two experimental batches. The pilot batches were used for the development of the experimental batches.

The preparation was carried out as follows:

- 1) Preparation of the melatonin mixture for the nucleus with volume excipients, gliding and lubricating excipients, binding excipients, for example bicalcium phosphate, lactose, aerosil, magnesium stearate, polyvinylpyrrolidon.

A non-limiting example of the granulate formulation is:

30	melatonin	2 mg
	mannitol	31 mg
	bicalcium phosphate	30 mg

polyvinylpyrrolidon	4.5 mg
aerosil 200	0.5 mg

2) Granulation

3) Calibration of the granulate

4) Addition of the retardant excipients, lubricants, volume and gliding excipients as, for example, hydroxypropylmethylcellulose, lactose, aerosil, magnesium stearate.

A non-limiting example of the unit formulation for the formation of the nucleuss is:

granulate (from above)	68 mg
hydroxypropylmethylcellulose	31 mg
lactose	75 mg
aerosil 200	0.35 mg
Mg stearate	1.65 mg

5) Formation of delayed release nucleus: controlled pressure compression in a tablet press to obtain nucleus hardness equal to 7-8 kN

6) Preparation of the melatonin solution for the "cortex" of the tablets, containing the following excipients: hydroxypropylmethylcellulose, lactose, ethyl alcohol, purified water.

A nonrestrictive example of formulation in solution for the "cortex" is:

melatonin	2.7%
hydroxypropylmethylcellulose	8.8%
lactose	6.4%
titanium bioxide	0.8%
ethyl alcohol	17.3%
purified water	64%

7) Application of the melatonin solution under pressure on the tablets, for the formation of the "cortex" testing for the quantity of melatonin by means of chemical analyses done on samples taken during manufacture.

Stages 2), 4), and 7) are essential for the preparation of the formulations which are the subject of this invention.

Example 2

Test in vitro for the release times of the active ingredients

The ready tablets, both in the pilot tests and in the experimental tests, as described above, were subjected to the following physical-chemical tests to obtain the following data:

- a) weight uniformity;
- b) hardness;
- c) friability;
- d) average content of melatonin;
- e) content uniformity;
- 6 f) disaggregation;
- g) time for the release of the active ingredient.

Weight uniformity, hardness, friability and disaggregation were checked by means of routine analyses according to standard procedures in the field. The average melatonin content and uniformity of content were determined by determination of the quantity of melatonin in the tablets.

Melatonin analysis

The pilot batch samples were analysed for their melatonin content by means of direct radioimmunoassay method (RIA), using an anti-melatonin antibody (Stockgrand LTD) and 2-^(125I)iodomelatonin (Amersham International) as a tracer. For every test various quantities of crushed tablets or supernatants from tablets incubated *in vitro* were used. RIA analysis was done in duplicate. Every sample was incubated overnight at 4° C together with the antibody and tracer.

The separation of the bound melatonin and the free melatonin was carried out by means of incubation with dextran-coated charcoal and subsequent centrifugation (3,000 g at 4°C for 20 mins.). The supernatant was quickly eliminated and the pellet was counted with a gamma - counter. The lower detection limit of the assay was 10 pg/ml.

The data were later confirmed by HPLC analyses on selected samples, in isocratic system.

Release times of the active ingredient.

The tablets prepared for the pilot tests were incubated in gastric/intestinal juices at 37° C on an oscillating tray and samples from the supernatants were taken out at predetermined times : 1, 5, 10, 30 minutes, 1, 2, 3, 4, 5, and 6 hours. In certain cases the incubation was carried out up to the eighth hour. The samples were analysed for the melatonin content as described above.

The experimental batches were produced on the basis of the data obtained from the pilot batches, and this was followed by the analysis of the content and by establishing the

average melatonin contents as previously described for the pilot tests. The determination of the release time was carried out in sixtuplicates for each of the experimental batches (Batch SP-01: 2 mg + 1 mg: Batch SP-02: 3 mg + 1 mg).

The data in vitro (analyses HPLC) are summarised in Tables I and II.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2

Table I

Release of melatonin from tablets 2 + 1 mg.

Time	10 min.	1 hour	2 hours	4 hours
Sample	Cortex (% of 1 mg)	Nucleus (% of 2 mg)	Nucleus (% of 2 mg)	Nucleus (% of 2 mg)
# 1	104	30	50	77
# 2	107	25	48	80
# 3	105	28	46	79
# 4	108	29	47	76
# 5	103	26	48	80
# 6	109	30	49	76
Average	106	28	48	78

Note: The data are expressed as percentages of the cumulative release.

Table II

Release of melatonin from tablets 3 + 1 mg.

Time	10 min.	1 hour	2 hours	4 hours
Samples	Cortex (% of 1 mg)	Nucleus (% of 3 mg)	Nucleus (% of 3 mg)	Nucleus (% of 3 mg)
# 1	95	24	48	87
# 2	98	22	49	83
# 3	96	27	47	84
# 4	97	25	50	85
# 5	99	27	49	86
# 6	97	25	47	85
Average	97	25	48	85

6 Note: The data are expressed as percentages of the cumulative release.

These results *in vitro* carried out with the experimental batches agreed with the results obtained in the pilot tests and show that the complete release of melatonin from the "cortex" took place within 10 minutes (95 – 106% of the quantity released); the release of the nucleus at 78 – 85% took place within four hours, 95% within the fifth hour from the start of incubation.

- 6 Laboratory analysis also confirmed weight uniformity; hardness and friability, in the standard set by us; the average melatonin content, homogeneous; the content and disaggregation, uniform.

On the basis of the data obtained, we proceeded to experimentation on healthy volunteers to determine the pharmacokinetics of melatonin.

EXAMPLE 3

PHARMACOKINETICS IN MAN (BLOOD AND SALIVA: RIA)

Materials and Methods

Subjects

Six adult subjects, 4 males and 2 females with an average age of 35 took part in the test. The volunteers had previously been informed of the purpose of the research and received a memorandum on how it would take place. All gave verbal consent. All were healthy and had never suffered from any chronic diseases; none of them had taken any medicines for the 2 weeks prior to the test nor during the period of research.

The subjects taking part in the test took one tablet of melatonin (2mg + 1 mg) at 8.45 a.m., in two groups of three subjects. The time of administration was chosen in so as not to interfere with the normal rhythm of melatonin secretion in the 12 hours afterwards. All the volunteers were exposed to light of an intensity of > 2000 Lux during the experiment.

- 24 In the first stage the determination of melatonin pharmacokinetics after oral administration was carried out by measuring melatonin levels in the plasma and in the saliva of 3 of the subjects for 12 hours following administration (considered as time "0").

Before the administration of melatonin, a sterile heparinized catheter was inserted in the antecubital vein of all the subjects. The obtained heparinized blood samples were immediately centrifuged and the plasma maintained at –20°C until it was analysed. The saliva samples were taken using special swabs (Salivette, Sarsdet, Germany). At the times indicated above, the volunteers chewed the swabs for about 1 minute. The saliva was later separated by centrifugation at 4°C and maintained at –20°C until it was assayed.

The blood and saliva samples were analyzed out simultaneously at times of 0, 30, 60, and 90 minutes and every hour for 12 hours starting from the second hour.

The volunteers were told not to drink coffee or use toothpaste for at least 30 minutes before every saliva sample.

In the second stage of the experiment, melatonin pharmacokinetics was determined in healthy volunteers, carried out by measuring melatonin levels only in the saliva, for 12 hours after administration (0 time). The saliva samples were taken as before at 0, 30, 60 and 90 minutes and every hour for 12 hours starting with the second hour.

Melatonin content in plasma and saliva.

All the plasma samples were analysed for their melatonin content by means of direct radioimmunoanalytic method (RIA), using an anti-melatonin antibody (Stockgrand Ltd.) and 2-^(125I)iodomelatonin (Amersham International) as tracer.

Two hundred micro-litres of plasma were used for every test. The analysis was carried out in duplicate. Every plasma sample was incubated over-night at 4°C with the antibody and tracer. The separation of the bound melatonin from the free melatonin was carried out by incubating with dextran-coated charcoal and then centrifuged (3000g at 4°C for 20 minutes). The supernatant was rapidly eliminated and the pellet counted with a gamma counter. The lower level of assessment is 10pg/ml.

Melatonin analysis in saliva was carried out using a direct radioimmunoanalytic method (RIA), using an anti-melatonin antibody (Stockgrand Ltd) and 2-^(125I)iodomelatonin (Amersham International) as tracer. Five hundred microlitres were analysed for every saliva sample; the test was duplicated. Every sample was incubated over-night at 4° C with the antibody and tracer.

The separation of the bound melatonin from the free melatonin was carried out with separation in a liquid phase. A second (anti-IgG antibody) and 6% PEG were added and then incubated for 4 hours at 4° C.

After centrifugation at 3000g per minute, the supernatant was quickly eliminated and the pellet counted with a gamma-counter. The lower limit of detection was 4 pg/ml. The cumulative results are shown in Figure 1.

The following conclusions are drawn from the data obtained:

- The melatonin content in the saliva faithfully reflects the pharmacokinetics in the peripheral blood. However, it should be remembered that maximum levels reached in

saliva are delayed in comparison with those obtained in blood. This phenomenon is clearly due to the transport of melatonin from the blood to the salivary glands and therefore in the saliva.

- The controlled release formula 1 mg + 2 mg is suitable for administration in man in that it achieves very low pharmacological levels (200 – 1500 pg/ml) for a period of 6 – 7 hours.
- The controlled release formula 1 mg + 2 mg results in levels of melatonin in the peripheral blood with a pattern very similar to the natural one (night-time melatonin).
- The controlled release formula 1 mg + 2 mg is suitable for use in sleep induction because levels of melatonin higher than 1000 pg/ml are obtained in the peripheral blood within 30 minutes from its administration.
- The controlled release formula 1 mg + 2 mg is suitable for the use in the maintenance of induced sleep because melatonin levels between 250 and 1000 pg/ml are maintained in the peripheral blood for about 6 hours after administration.

Comparable data from the kinetics viewpoint were obtained with the administration of melatonin in the formula 1 mg + 3 mg, clearly proving that the invention can be used with different quantities of the active compound.

Example 4

EFFECTS ON SLEEP IN MAN

Materials and Methods

Subjects and treatment

Ten patients (average age 56 ± 3.6 years) suffering from psychophysiological insomnia, were used in the study. All of were classified according to The International Classification of Sleep Disorders, Revised Edition, ASDA, 1997. All were informed and gave their verbal consent to take part in the study. All received case report forms to fill in the following parameters: Total Sleep Time (TST), Sleep Latency (SL), Wake After Sleep Onset (WASO), Number of Awakenings (NA).

The treatment was in "crossover" and lasted a week for every formulation. Prior/after (according to a random assignment) to the treatment with controlled release formulations containing melatonin according to this research (1 + 2 mg of melatonin), the subjects took standard formulations of melatonin in 3 mg tablets, followed by a washout period of three

days. In both treatments the tablets were given at 22.30 hours. Then, according to the data gathered, sleep effectiveness was calculated.

Results

The results are summarised in Table III.

TST, SL and WASO are expressed in minutes. NA is the numerical value of the awakenings. SE is the calculated percentage of time spent asleep minus the time spent awake after beginning of sleep.

TABLE III

Parameter	Baseline	Melatonin Normal	Melatonin Finding
TST	270 ± 24	290 ± 18 ^a	330 ± 34 ^{a,b}
SL	55 ± 7.5	38 ± 5.6 ^a	34 ± 8.3 ^{a,b}
WASO	42.2 ± 23.1	44.2 ± 19.4	27.3 ± 10.2 ^{a,b}
NA	6.2 ± 2.4	7.5 ± 2.7	4.3 ± 1.6 ^{a,b}
SE (%)	84.3	84.7	91.7 ^{a,b}

a = significantly different from the Baseline.

b = significantly different from normal Melatonin

The above data clearly show that the standard formulations of melatonin, while diminishing sleep latency (SL) and marginally influencing TST, do not have any significant effects on SE, because they do not reduce WASO and NA.

On the contrary, the controlled-release formulations containing melatonin according to this invention, in the same dosage, improved all sleep parameters : TST, SL, WASO, NA and consequently SE.

The data on WASO and NA are to be considered extremely important within the embodiment of this invention for their importance on sleep quality, evidently due to the time-controlled release effect of melatonin.

This invention has the following advantages compared with solutions used until now: